

GEOGRAPHIC VARIATION OF CHLOROPLAST DNA HAPLOTYPES IN *Acacia aulacocarpa* A. Cunn. ex Benth

Anthonius YPBC Widyatmoko¹ and Susumu Shiraishi²

Received : 29 January 2013, Accepted : 26 June 2013

ABSTRACT

The geographic variation of chloroplast DNA (cpDNA) haplotypes of *Acacia aulacocarpa* was investigated among 18 natural populations. These populations represent the geographical range of the species in New Guinea Island and Queensland. Single strand conformation polymorphism (SSCP) was used for the analysis. Two non-coding regions of cpDNA, the intron region of the *trnL* gene and the intergenic spacer region between the *trnP* and *trnW* genes, were analyzed, and four haplotypes (A, B, C, and D) were recognized. The haplotype distribution corresponded with the geographic distribution of the populations. Based on four cpDNA haplotypes, the eighteen populations were divided into three groups: New Guinea Island, Northern Queensland and Southern Queensland. Haplotype C was observed only in the New Guinean populations, while the other three haplotypes (A, B, and D) were found in Queensland only. All of these three haplotypes were observed in Southern Queensland, whereas haplotype B was found only in the Northern Queensland populations. The cpDNA haplotype diversity of this species seemed to be highest in southern Queensland.

Keywords: *Acacia aulacocarpa*, geographic variation, Chloroplast DNA, non-coding region, haplotype, PCR-SSCP

I. INTRODUCTION

The genus *Acacia* is a common forest tree species originated from Papua and Papua New Guinea (New Guinea Island), Australia and neighboring islands. Among *Acacia* species, *A. mangium* Willd. and *A. auriculiformis* Cunn. ex Benth. are considered as the most important species for plantation, and some genetic improvement programs have been carried out in Southeast Asia (Thompson, 1994). In addition, *A. aulacocarpa* A. Cunn. ex Benth and *A. crassicarpa* Cunn. ex Benth are widely distributed in tropical areas. These species are utilized in forest plantation for pulp and paper. Genetic diversities in *A. mangium*, *A. auriculiformis*, and *A. crassicarpa* have been discussed by Moran *et al.* (1988 and 1989), Khasa *et al.*, (1994), Wickneswari and

Norwati (1993), Butcher *et al.*, (1998), Widyatmoko *et al.*, (2010) and Widyatmoko and Shiraishi (2011). Several studies have been carried out in species trial, provenance trial, nursery practice and fertilization to determine superior growth performance, adaptability, pest and disease resistance of the genus (Thompson, 1994). Phylogenetic study on *Acacia* has been reported by Brown *et al.* (2008), Byrne *et al.* (2001 and 2002) and Clarke *et al.* (2000).

A. aulacocarpa belongs to the section of *Juliflorae*, subgenus *Phyllodineae* (Brain and Maslin, 1996; Pettigrew and Watson, 1975). This species is naturally distributed in diverse habitats over a wide geographic range, including four states in northern and eastern Australia (Queensland/QLD, New South Wales/NSW, Northern Territory/NT, and Western Australia/WA), Papua New Guinea (Western Province/WP), and Indonesia (Papua/PI). Based on morphological differences, this species has been divided into 5 subspecies: *A. aulacocarpa* subsp. A (NT, WA), *A. aulacocarpa* subsp. B (WP, PI), *A. aulacocarpa* subsp.

¹Center for Forest Biotechnology and Tree Improvement, Yogyakarta

²Laboratory of Silviculture, Faculty of Agriculture, Kyushu University, Fukuoka, Japan

Corresponding Author: aviwicaksono@yahoo.com

C (northern QLD), *A. aulacocarpa* subsp. D (northern QLD), and *A. aulacocarpa* subsp. E (southern QLD, NSW) (Thomson, 1994). McDonald and Maslin (2000) revised taxonomic of *Acacia aulacocarpa* Cunn. ex Benth. and its seven close relatives, mainly based on their mode of pod dehiscence. These species divided into two subgroups: *A. aulacocarpa* subgroup and *A. crassicarpa* subgroup. *Acacia aulacocarpa* subgroup consists of *A. aulacocarpa*, *A. celsa* and *A. disparrima*, while *A. crassicarpa* subgroup comprise of *A. crassicarpa*, *A. lamprocarpa*, *A. midgleyi*, *A. peregrina* and *A. wetarensis*.

Field trials regarding to the species have been carried out in several countries. *A. aulacocarpa* subspecies C (Thomson, 1994), is known as highly productive and high survival rate at most of the trial sites. This species demonstrates high pulp productivity (330 kg/m³) especially for kraft pulping. In comparison with other *Acacia*'s species (*A. auriculiformis*, *A. cincinnata*, *A. crassicarpa* and *A. mangium*), *A. aulacocarpa* yields more for pulping (Balodis and Clark, 1998).

Single-strand conformation polymorphism (SSCP) is a simple, fast and very sensitive method for the identification of nucleotide-sequence polymorphisms. This technique is performed without sequencing and able to detect small changes in DNA such as single base substitutions, deletions, and insertions (Orita *et al.*, 1989). Hayashi (1991) combined this method with PCR, and developed a simple method of analysis referred to as PCR-SSCP. This method has greatly increased the efficiency of the marker and appears to be a useful tool for studying genetic variation. It has been utilized in studies on the detection of inheritance and molecular variation (Bodenes *et al.*, 1996), chloroplast DNA variation, typing and identification (Maeda and Shiraishi, 1997; Watano *et al.*, 1995), and genetic mapping (Judelson *et al.*, 1995). Hongyo *et al.* (1993) reported the improvement of SSCP method becomes simple method called 'Cold-SSCP'. Recently, the SSCP method has been further modified and utilized for DNA genome analysis (Grace *et al.*, 1995).

The chloroplast genome is conservative in its evolution. Chloroplast DNA (cpDNA) possesses

many peculiar characteristics, such as uniparental inheritance, the lack of recombination and repeated sequences, and low variability compared with nuclear DNA (Birky, 1988; Wolfe *et al.*, 1987). These characteristics lead to the expectation that diversity within a population is very low and differentiation between populations might be easily detected. This would be especially observable in species with geographically isolated populations and/or small populations.

Studies of the genetic diversity of the *Acacia aulacocarpa* have been reported by Widyatmoko and Shiraishi (2010). In this study, however, only six samples have been used and compared to the others three *Acacia* species of which *A. aulacocarpa* had the highest genetic diversity. Information of genetic diversity for all natural distribution of *A. aulacocarpa* is needed. Therefore, research was carried out with main objective to elucidate genetic diversity and genetic distribution of the species.

II. MATERIAL AND METHOD

A. Plant Materials

A total of 87 individuals (seeds) originating from 18 populations were used in this study (Figure 1 and Table 1). The seed samples were obtained on 1994 from the Australian Tree Seed Centre of the CSIRO (Commonwealth Scientific and Industrial Research Organization), Australia and from the Forest Tree Improvement Research and Development Institute, Indonesia. The populations were distributed in Indonesia, Papua New Guinea, and QLD (corresponding populations: 1, 8 and 9, respectively). Population latitudes ranged from 7.45°S (Papua, Indonesia) to 27.17°S (Samford, QLD); altitudes ranged from 6 m a.s.l (N. Yeppoon, QLD) up to 720 m a.s.l (Buckley, QLD). For each population, 2-6 trees are represented; only one seed from each individual tree (n=87) was used. Similar number of samples per population for analysing chloroplast DNA variation also reported by Zufahmi *et al.* (2010) and Bordacs *et al.* (2002).




Figure 1. Natural Distribution of *A. aulacocarpa* and location of population invested
Remarks: Population number refers to Table I.

B. DNA Extraction

Total genomic DNA was extracted using a modified SDS method (Guan *et al.*, 2011). Each seed was grounded using 400 μ l SDS extraction buffer, which contained of 50 mM Tris-HCl (pH 9.0), 1% (w/v) SDS, 10 mM EDTA, and 0.5% (v/v) 2-Mercaptoethanol. After incubation at 65°C for 60 min, 200 μ l of 7.5 M ammonium acetate was added. The solution was kept on ice for 30 min, and was then centrifuged at 0°C at 15,000 rpm for 40 min. The sample in aqueous phase (400 μ l), was transferred to a new tube, and the DNA was precipitated by the addition of 400 μ l isopropanol. After circa 10 min, the precipitate was collected by centrifugation at 15,000 rpm for 10 min. The supernatant was completely removed, and the pellet was washed twice with 1.0 ml of 70% Ethanol. After washing, the pellet was then vacuumed in an evaporator for 2 min, then it was resuspended in 100 μ l purified H₂O. Finally, the crude solution was purified using a GeneClean III Kit (BIO 101).

C. PCR Amplification

The 10 μ l PCR reaction mixture contained 2 ng genomic DNA, 0.25 μ M of each primer, 10 mM Tris-HCl (pH 8.3), 10 mM KCl, 3.0 mM MgCl₂, 0.2 mM each of dNTPs, and 0.5 unit of AmpliTaq DNA polymerase, Stoffel Fragment (Perkin-Elmer). PCR was performed using a GeneAmp PCR System, Model 9600 (Perkin-Elmer). The conditions of amplification were as follows: 95°C for 60 s, and 30 cycles of 30 s at 94°C, 30 s at 55°C, and 90 s at 72°C, followed by 60 s at 72°C.

In this study, six pairs of primers (Table 2) were used to amplify six cpDNA regions. The four regions were as follows: one intron region of the *trnL* gene (CS2; B49317/A49855) and three intergenic spacer regions, namely, those between the *trnL* and *trnF* genes (CS3; B49873/A50272), between the *trnD* and *trnY* genes (CS4; S4U/CS4L), and between the *trnP* and *trnW* genes (CS5; CS5U/CS5L). Two regions (CS5a and CS5b) originated from the CS5 region were also amplified with two primer pairs (CS5aCS5aU/CS5aL and CS5bU/CS5bL), respectively.

Table 1. Details of seedlot number and location of sample populations of *Acacia anulocarpa*

No.	Population	No. of seeds	Seedlot No.	Latitude (°S)	Longitude (°E)	Altitude (m a.s.l.)
New Guinea						
1.	Kuel, Irian Jaya	6	1001, 1002, 1003, 1004, 1005, 1006	7.45	140.45	45
2.	Makapa WP	5	16974-M (3, 7, 12, 16, 19)	7.56	142.35	35
3.	Isago Arimia River WP	5	16948-AI (3, 4, 5, 6, 7)	8.10	142.41	10
4.	Duaba WP	5	16949-AD (2, 4, 6, 8, 10)	8.13	142.58	25
5.	Wasua Pedeya WP	5	16950-AW (1, 11, 21, 31, 37)	8.17	142.52	52
6.	Kapal NW of Wipim WP	5	16981-BVG (1651, 1653, 1654, 1655, 1659)	8.37	142.47	40
7.	Derideri E Morehead WP	5	16989-MM (969, 970, 971, 974, 975)	8.40	141.50	30
8.	Wipim WP	5	16976-KER (2, 6, 10, 13, 16)	8.47	142.52	45
9.	Balimo District	5	16946-AKO (1, 12, 22, 32, 42)	8.50	142.58	12
Northern Queensland						
10.	Old Rock Hart Airstrip	5	18358-GJM (1377, 1383, 1384, 1385, 1386)	12.50	143.18	15
11.	Mcil Wraith Range	4	18001-T (1142, 1143, 1145, 1146)	13.44	143.20	500
12.	Piccaninny Creek	2	16136-JMO (1459, 1463)	13.90	142.48	40
13.	10 K NW Mt. Molloy	5	17905-TREE (1, 2, 3, 4, 5)	16.40	142.15	420
14.	Garioch	5	13866-BVG (2939, 3940, 2941, 2942, 2943)	16.40	145.18	400
15.	Buckley Logging Area	5	13865-BVG (2921, 2935, 2936, 2937, 2938)	17.90	145.37	720
Southern Queensland						
16.	3K S Mt. Larcom	5	17739-GB (22, 25, 27, 28, 73)	23.50	151.00	70
17.	N of Yeppon	5	14591-PWO (174, 175, 178, 184, 185)	23.60	150.45	6
18.	Samford	5	17891-GB (96, 97, 98, 99, 100)	27.17	152.51	50

Table 2. Primer sequences of cpDNA non coding regions used in this study

Region	Code	Primer sequence (5' – 3')	Reference	
<i>trnL</i> intron	CS2	B49317	CGAAATCGGTAGACGCTACG	Taberlet <i>et al.</i> (1991)
		A49855	GGGGATAGAGGGACTTGAAC	
		sCS2U	tgtaaaacgacggccagtCGAAATCGGTAGACGCTACG	
		sCS2L	caggaaacagctatgaccGGGGATAGAGGGACTTGAAC	
<i>trnL</i> - <i>trnF</i>	CS3	B49873	GGTTCAAGTCCCTCTATCCC	Taberlet <i>et al.</i> (1991)
		A50272	ATTTGAACTGGTGACACGAG	
<i>trnD</i> - <i>trnY</i>	CS4	CS4U	TGACAGGGCGGTACTCTAAC	Shiraishi <i>et al.</i> (unpublished)
		CS4L	CGATGCCCGAGTGGTTAATG	
<i>trnP</i> - <i>trnW</i>	CS5	CS5U	TTGGTAGCGTGTGTTGTTTGGG	Isoda and Shiraishi (1999)
		CS5L	TACGGCATCAGGTTTTGGAGAC	
	CS5a	CS5aU	GTGACCCCAATTCATCTTGC	Newly-designed In this study
		CS5aL	TTCCCTAGTTAACTCTGGCT	
	CS5b	CS5bU	TAAATCATTTTATGTGACCC	Newly-designed In this study
CS5bL	CGTCATGTCCCTATGGCT			

Lower case indicates -21M13 primer or M13 Rev primer (Perkin-Elmer)

D. SSCP Analysis

PCR products (2.5 µl) were mixed with 12 µl of loading buffer containing 77% formamide, 0.25% BPB, and 1x TBE. The solution was concentrated for 15 minutes using a vacuum evaporator, denatured for 5 min at 94°C, then immediately cooled on ice, and loaded onto 7% polyacrylamide (39:1 acrylamide to bis acrylamide) TBE gel. Depending on the length of PCR product, the electrophoresis was run for 2 or 4 hours at 1.5 V/cm and at a constant temperature of 10°C in 1x TBE running buffer. The gel was stained with a 0.5 µg/ml solution of ethidium bromide in 1x TBE buffer for 15 min. The SSCP bands were visualized using a 302 nm UV transilluminator.

E. Cloning of DNA Fragments

The PCR and SSCP assays revealed two duplicated DNA fragments in the CS5 region (Fig. 3). To determine the sequence of each fragment, a cloning technique was adopted. Each fragment was cut from the gel and used as a template for the next PCR. PCR amplification was performed again for each fragment using the same primers as those used for CS5 (CS5U and CS5L). The two PCR products were ligated with pGEM vector (Primega), and were transformed into *E. coli* (JM109 High Efficiency Competent Cell) according to the method recommended by the

suppliers. The inserted DNA regions were directly amplified from colonies using colony PCR with the following primers: 5' TCCGGCTCGTATGT-TGTGTGGA-3' and 5'-GTGCTGC-AAGGCGATTAAGTTGG-3' (Hattori, 1997). DNA amplification was carried out using the above-mentioned PCR composition, except with an extremely small *E. coli* colony as a template DNA. PCR was performed using a GeneAmp PCR System, Model 9600 (Perkin-Elmer) with the thermal profile of 95°C for 60 s; 29 cycles of 15 s at 96°C, and 120 s at 70°C; followed by one cycle of 15 s at 96°C and 10 min at 70°C. The PCR products were purified with a GeneClean III Kit (BIO 101) and used as templates for sequencing.

G. Sequencing

In addition to the colony-PCR products mentioned above, the CS2 region was amplified as templates for sequencing using a primer pair (sCS2U/sCS2L) shown in Table 1. A 20 µl reaction mixture contained 4 ng of genomic DNA, 0.25 µM of each primers, 20 mM Tris-HCl (pH 8.0), 100mM KCl, 2.0 mM MgCl₂, 1 mM DTT, 0.25 mM of each dNTP, and 1.0 unit of *TaKaRa ExTaq*™ (TAKARA). PCR was performed with a Gene Amp PCR System, Model 9600 (Perkin-Elmer). The conditions of amplification were 95°C for 60 s, and 30 cycles of

30 s at 94°C, 30 s at 45°C, and 90 s at 72°C, followed by 60 s at 72°C. In order to separate the amplified DNA from residual primers, all reactions were pooled and subjected to electrophoresis in a 1.5% (w/v) low-melting-point agarose gel using TBE buffer. The desired DNA fragment was cut out of the gel and recovery was obtained using QIAEX II (QIAGEN). Sequencing was performed with a Thermo Sequanase pre-mixed cycle sequence kit (Amersham) and two sequence primers (Perkin-Elmer -21M13: 5'-TG TAAAACGA-CGGCCAGT-3' and M13 Rev: 5'-CAGGAAACAGCTATGACC-3') using HITACHI SQ5500 Sequencer.

III. RESULT

A. PCR-SSCP Profile

Of the four DNA regions where the cpDNA variation was surveyed in *A. aulacocarpa*, two regions (CS2 and CS5) showed polymorphisms, and the other two regions (CS3 and CS4) were monomorphic. In the SSCP profile of the CS2

region, two bands appeared that corresponded to the two single-strand DNAs (ssDNA) of the PCR product. Figure 2 depicts three SSCP types showing different migration patterns. These three types were referred to as $cs2^1$, $cs2^2$ and $cs2^3$. In the NGI populations, all of the samples were of the $cs2^2$ type (lanes b, d, f, h, i, and k in Figure 2). In contrast, 3 different types were observed in the QLD populations. The $cs2^1$ type was recognized in the populations of N. Yeppoon and 3K S Mt. Larcom (lanes e and g), while $cs2^3$ was found only in the one sample from Samford (lane j). The remaining samples from QLD (lanes a and c) were of the same type ($cs2^2$) as those from NGI. Within-population polymorphism was detected in only two populations, namely, those from Samford and N. Yeppoon.

In the CS5 regions, three SSCP types were observed and all samples possessed four fragments (Fig. 3B). As each fragment and a single-strand DNA were in a one-to-one correspondence, the discovery of these four fragments indicates that two DNA regions were amplified by the PCR with primers CS5U and CS5L.




Figure 2. Cold SSCP profiles of chloroplast *trnL* intron (CS2) in *Acacia aulacocarpa*

Remarks: Fragment length of CS2 is 578 bp (same as in Fig. 6)

a-k, individual tree number: a, 13965-BVG 02921; b, 16946-AKO 00001; c, 13966-BH 02939; d, 16947-M 0000003; e, 14591-PWO 00174; f, 16948-AI 000003; g, 17739-GB 000022; h, 16949-AD 000002; i, 1001; j, 17891-GB 000096; k, 16950-AW 000001.

$cs2^1$, "fast" allele; $cs2^2$, "medium" allele; $cs2^3$, "slow" allele

B. Polymorphism in CS5 Region

In the SSCP profile of the CS5 regions (Fig. 3B), all of the samples possessed four bands. These findings clearly suggest a duplication of the region between the *trnP* and *trnW* genes. Three types of band patterns were observed in this

region. The lanes a, b, and c, which showed slower migration patterns, belonged to the NGI populations. On the other hand, lanes e and f, which showed faster migration, were from the QLD populations. Furthermore, intermediate migration was recognized in a sample (lane d) from the Samford population (QLD).




Figure 3 PCR product (A) and Cold SSCP (B) profiles of chloroplast *trnP-trnW* spacer (CS5) in *A. aulacocarpa*

Remarks: A, B, and C, three different PCR products.

a-f, individual tree number: a, 1001; b, 16946-AKO 00001; c, 16947-M 0000003; d, 17891-GB 000096; e, 17905-TREE 1; f, 18358-GJM 1377; M, 100 bps ladder

The agarose gel electrophoretic profile of the PCR products (Figure 3A) revealed that each sample possessed 2 bands in which the fastest band (A) was thicker than the slower bands (B and C). The sequence of the duplicated regions was clarified by cloning three DNA fragments (A, B, and C). Figure 4 depicts the sequences of fragments A, B, and C. These three sequences were highly homologous, especially fragments A and B, which were completely identical except for an insertion/deletion (indel) of 40 bps. The sequence of fragment C was different from the former two sequences as regards five of the indels and four substitutions. From the sequence homology, these three sequences (A, B, and C) could be grouped into CS5a and CS5b. The fragments A and B reflected the CS5a region, and the fragment C corresponded to CS5b. Based on

the sequence information, two primer pairs were newly designed to amplify the two regions (CS5a and CS5b) specifically, and SSCP variation was investigated in each region (Figure 5A and 5B).

In the CS5a region, two SSCP types with different migrations were detected (Figure 5A). The two types were referred to as $cs5a^1$ and $cs5a^2$, respectively. The $cs5a^2$ was found in only one sample from the Samford population (lane a). On the other hand, in the CS5b region, two SSCP types with different migrations were observed (Figure 5b). The two types were referred to as $cs5b^1$ and $cs5b^2$. The odd lanes in Figure 5B were the samples from Queensland populations, whereas the even lanes were from NGL. This revealed that these two groups, NGL and Queensland, possessed quite different SSCP types of CS5b region.




Figure 4. A partial Sequences of chloroplast *trnP-trnW* (Cs5) spacer of the three different PCR products (A, B, and C) as shown in Figure 3

Remarks: Two primer pairs (CS5a and CS5b) were designed to amplify the two duplicated regions. A dash (-) denotes a deletion




Figure 5 (A). Cold SSCP profiles of CS5a regions in *Acacia auilacocarpa* a-k, individual tree number : a, 17891-GB 000096; b, 17739-GB 000073; c, 1006, d, 16950-AW 000037; e, 17891-GB 000100; f, 16971-KER 00016; g, 17905-TREE 5; h, 16981-BVG 01659; i, 18358-GJM 1386; j, 16989-MM 000975; k, 16136-JMO 01463. $cs5a^1$, "fast" allele; $cs5a^2$, "slow" allele

(B). Cold SSCP profiles of Cs5b regions in *Acacia auilacocarpa* a-k, individual tree number : a, 17905-TREE 1; b, 16976-KER 00002; c, 18358-GJM 1377; d, 16981-BVG 01651; e, 1800-T 1142; f, 16989-MM 000969; g, 16136-JMO 01459; h, 16946-AKO 00012; i, 13865-BVG 02935; j, 16947-M 0000007; k, 13866-BVG 02940. $cs5b^1$, "fast" allele; $cs5b^2$, "slow" allele

C. Sequences of Polymorphic Regions

The sequences of SSCP types in each polymorphic region are shown in Figure 6. It was confirmed that all of the SSCP polymorphisms in the three regions reflected differences in sequence. In the cpDNA *trnL* intron region (CS2) in which three SSCP types were detected, all sequence types had the same length (578 bps); furthermore, no indel was detected. Single-base substitution, however, was recognized among the three types. $cs2^1$, observed in 3K S Mt. Larcom and N. Yeppoon populations reflected a single-base transition (G to A) in the 177th base. However, the $cs2^3$ detected in one sample in the Samford population reflected a single transition (C to T) in the 228th position.

According to the cpDNA intergenic spacer between *trnP* and *trnW* (CS5), two regions were analyzed using the different primer pairs. In CS5a, the two types were distinguished by a 5-base

duplication and a single-base transversion (A to C). In CS5b, which clearly separated NGI from QLD, the sequence length differed between the two types. In this region, a tandem repeat with a 20-bp core sequence was recognized. $cs5b^2$ appeared only in the NGI populations that had a twice-repeat sequence, whereas $cs5b^1$ was detected only in QLD populations that had a four-fold-repeat. Moreover, a single-base substitution was detected between $cs5b^1$ and $cs5b^2$.

D. Chloroplast DNA haplotype Distribution

Four different cpDNA haplotypes have been determined based on the combination of SSCP types in the three regions (Table 3). Haplotype A was a combination of $cs2^1$ in CS2 region and $cs5a^1$ and $cs5b^1$ in CS5. Haplotype B was a combination of the $cs2^2$, $cs5a^1$, and $cs5b^1$. Haplotype C consisted of $cs2^2$, $cs5a^1$, and $cs5b^2$, and haplotype D was a combination of $cs2^3$, $cs5a^2$, and $cs5b^1$.

```

                                                    60
GACTTAATTG GATTGAGCCT TGGTATGGAA ACCTACCAAG TGATAACTTT CAAATTCAGA
.....
.....

51                                                    120
GAAACCTGG AATTAACAAT GGGCAATCCT GAGCCAAATC CTGTTTCCG AAAACCAAGA
.....
.....

121                                                    180
AGAGTTCAGA AAGGGAGAAT AAAATAAAAA AGGATAGGTG CAGAGACTCA ACGGAAGCTG
.....A...
.....

181                                                    240
TTCTAACAAA TGGAGTTGAC GACATTTTCGT TTCCTTAGTA AGGATTTTCGT TTCGTTAGTA
.....T...
.....

241                                                    300
AAGAGATCCT TCCATCGAAA CTCCAGAAAA GAAAGGATCA AGGATGAGCA TACGTATACG
.....
.....

301                                                    360
TACTGAAATA CTATTTCAAT TGATTAGACC AGACAGACCC AAAATCTCTA TTTTTATTT
.....
.....

361                                                    420
TAATATTTAT ATGACAAATA AAAGATGTGA ATCGATTCCA AGTTGAAGAA AGAATCAAAT
.....
.....

421                                                    480
ATTAATTGAT CAAATTCACT CCATGATAGT CTGATAGATT TTTTGAAGAA CTGATTAATC
.....
.....

481                                                    540
AGACGAGAAT AAAGATAGAG TCCATTCTA CATGTCAATA CCGACAATAA TGAAATTTAT
.....
.....

541                                                    578
AGTAAGAGGA AAATCCGTCG ACTTTAGAAA TCGTGAGG
.....
.....

```

Figure 6. Sequences of chloroplast *trnL* intron (CS2) in *A. aulacocarpa*

Remarks : Only positions different from the upper sequence are shown. Each line corresponds to the different allele. The upper line, *cs*² allele; the middle line, *cs*^{2'} allele; the lower line, *cs*^{2'} allele

Table 3. Allele combination of each chloroplast DNA haplotype

CpDNA haplotypes	Regions		
	CS2*	CS5a**	CS5b**
A	1	1	1
B	2	1	1
C	2	1	2
D	3	2	1

* Chloroplast *trnL* intron

** Chloroplast intergenic spacer between *trnP* and *trnW*

IV. DISCUSSION

A. SSCP and Sequence Polymorphism

Recently, PCR-SSCP analysis has proved to be a very sensitive, rapid, and easy method for detecting point mutations and other DNA polymorphisms (Rehben *et al.*, 1997; Watano *et al.*, 1995; Bodenes *et al.*, 1996; Watanabe *et al.*, 1997; Maeda and Shiraishi, 1997). So far, extensive use of this technique has been limited by disadvantages such as the need for radioactive labeling, time-consuming electrophoresis, and an accurate DNA concentration. These factors play an important role in the sensitivity and resolution of SSCP. A new method referred to as "Cold-SSCP" (Hongyo *et al.*, 1993) enabled SSCP profiles to be visualized directly with ethidium bromide staining.

The effectiveness of PCR-SSCP for detecting cpDNA haplotype polymorphism in *A. aulacocarpa* was demonstrated in this study. Orita *et al.* (1989) and Spinardi *et al.* (1991) have reported differences in SSCP band migration caused by single-base substitutions. In the present study, this method was adopted to enable single-base substitution and indel detection. In both CS5 regions (CS5a and Cs5b), sequence length variations and single-base substitutions were detected as mobility shifts in both SSCP bands. In the CS2 region, SSCP was able to distinguish three sequences that differed from each other by a single-base substitution.

B. Geographical Variation in Chloroplast DNA

In spite of the conservative evolution of the chloroplast genome, intraspecific variations of cpDNA have been widely reported (Levy *et al.*, 1996) and cpDNA has become increasingly valuable as a genetic marker to detect intraspecific variations. CpDNA markers have been used to examine population genetics and structure, haplotype diversity, genetic diversity, and breeding systems (Hong *et al.*, 1993; Levy *et al.*, 1996; Watano *et al.*, 1995). In the present study, cpDNA was adopted to estimate the geographic variation of *A. aulacocarpa*. Widyatmoko and Shiraishi (2011) reported cpDNA variation of 6 samples of *A. aulacocarpa* using intergenic spacer between the *trnP* and *trnW* gene. The NGI and QLD tend to be separated based on the cpDNA haplotypes.

In this study, cpDNA haplotype variations in *A. aulacocarpa* in the NGI and the QLD were detected in two non-coding regions (the *trnL* intron and the intergenic spacer between the *trnP* and *trnW* genes) by using PCR-SSCP.

Two duplicated regions were detected in the region between the *trnP* and *trnW* genes (CS5). By cloning and sequencing the DNA fragments, two new pairs of primers were designed, thus enabling the two regions to be specifically amplified by PCR. One of these two regions, maybe CS5b, seems to be a pseudogene. Highly homologous arrangements with cpDNA are also observed in nuclear DNA (ncDNA) (Birky, 1988). If the region was located in ncDNA, the heterozygous type with *cs5b*¹ and *cs5b*² should have appeared. However, no heterozygote was detected in any of the 87 samples analyzed in this study. Therefore, both regions were presumed to be located on the cpDNA. However, it cannot be ruled out that one of the two regions is located in a genome other than the cpDNA genome. A total of 7 SSCP types were detected in the three regions, and 4 cpDNA haplotypes were identified based on a combination of the 7 SSCP types (Table 2).

Table 3 shows the relative frequencies of the four cpDNA haplotypes in 18 populations. Based on the distribution of the four identified cpDNA haplotypes, these populations could be divided into three groups. One group consisted of nine populations that were located in NGI. This group could be characterized by the appearance of haplotype C, which this haplotype appeared only in NGI populations, and was not observed in the other populations. Another group of six populations in northern QLD was located at the latitude between 13 - 18°S. This group was characterized by haplotype B. The haplotype B appeared also in southern QLD populations. The third group, consisting of three populations, was distributed in southern QLD (at a latitude between 23 - 28°S). Unlike the other two groups, this group included three haplotypes, excluding haplotype B. Furthermore, variation in haplotype could be recognized between populations as well as within populations. These three populations can be divided into two more subordinate, smaller groups based on haplotype frequencies. Two populations, 3K S Mt. Larcom and N of Yeppoon, were distributed at an altitude of

Table 4. Distribution of chloroplast DNA haplotypes of each *Acacia aulacocarpa* population in New Guinea and Queensland

No	Population	No. of seeds	cpDNA haplotype (%)			
			A	B	C	D
New Guinea						
1.	Kuel, Irian Jaya	6	0.000	0.000	1.000	0.000
2.	Makapa WP	5	0.000	0.000	1.000	0.000
3.	Isago Arimia River WP	5	0.000	0.000	1.000	0.000
4.	Duaba WP	5	0.000	0.000	1.000	0.000
5.	Wasua Pedeya WP	5	0.000	0.000	1.000	0.000
6.	Kapal NW of Wipim WP	5	0.000	0.000	1.000	0.000
7.	Derideri E Morehead WP	5	0.000	0.000	1.000	0.000
8.	Wipim WP	5	0.000	0.000	1.000	0.000
9.	Balimo District	5	0.000	0.000	1.000	0.000
Northern Queensland						
10.	Old Rock Hart Airstrip	5	0.000	1.000	0.000	0.000
11.	Mcil Wraith Range	4	0.000	1.000	0.000	0.000
12.	Piccaninny Creek	2	0.000	1.000	0.000	0.000
13.	10 K NW Mt. Molloy	5	0.000	1.000	0.000	0.000
14.	Garioch	5	0.000	1.000	0.000	0.000
15.	Buckley Logging Area	5	0.000	1.000	0.000	0.000
Southern Queensland						
16.	3K S Mt. Larcom	5	1.000	0.000	0.000	0.000
17.	N of Yeppon	5	0.800	0.200	0.000	0.000
18.	Samford	5	0.000	0.800	0.000	0.200
			0.104	0.414	0.471	0.011

Frequencies of cpDNA haplotypes are shown in parentheses

around 23°S; these two populations had common haplotypes. The other population, namely, the Samford population, shared a common haplotype with the northern QLD group.

In *A. mangium*, a correlation between the cluster analysis and geographical distribution has been reported (Butcher, 1998). *A. mangium* has been grouped into four distinct regions, the Sidei and Ceram Islands, NGI, the northern part of northern QLD, and the southern part of northern QLD. The group in the northern part of northern QLD was more closely related to the NGI group than to that of the southern part of northern QLD. Similar results have been reported in *A. auriculiformis* (Wickneswari and Norwati, 1993). This species was divided into three distinct groups, the Northern Territory (NT), QLD, and NGI; the populations of QLD were closely

related to those of NGI. In the present study, however, a strong separation was observed between the NGI and QLD populations as regards the cpDNA variation. Since Australia and NGI were separated by the rising sea level during the postglacial period ca. 10,000 years ago, it might be easily predicted that the species adapted to different environments and that the gene flow between Australia and NGI was obstructed.

A rare haplotype was detected in a restricted population. Such a phenomenon has also been recognized in the isozymes of *A. mangium*, *A. auriculiformis*, and *A. crassiparva* (Moran *et al.*, 1989; Wickneswari and Norwati, 1993; Khasa *et al.*, 1994). Rare allelic variants at a particular locus were detected in only one or two populations; this could be due to mutations that arose during the postglacial period. Compared with *A. mangium*, *A.*

auriculiformis, and *A. crassicarpa*, *A. aulacocarpa* is more widely distributed and formed larger forests in QLD. Hence, a relatively high level of genetic diversity was expected in this territory; evidence for this hypothesis was given by the polymorphic cpDNA haplotype.

The clustering of the *A. aulacocarpa* population in NGI and QLD based on cpDNA haplotypes was in agreement with that based on the morphological characteristics. No population of NT was investigated in this study. The NGI group appears to belong to *A. aulacocarpa* subsp. B. The northern QLD group is presumed to correspond to subsps. C and D, whereas the southern QLD group corresponds to subsp. E. If these correspondences are accurate, the differentiation of cpDNA haplotypes in this study supports the subspecies classified by Thompson (1994). Distribution of eight species those clustered into 2 groups (McDonald and Maslin, 2000) also related with cpDNA haplotypes. *A. aulacocarpa* group (*A. aulacocarpa*, *A. celsa* and *A. disparrima*) distributed in northern and southern QLD. However, *A. crassicarpa* group (*A. crassicarpa*, *A. midgleyi*, *A. peregrina*), distributed in NGI and northern QLD. Thus, both groups can be separated with cpDNA haplotypes, except some population of *A. crassicarpa* and *A. midgleyi* those distribute in southern QLD. No samples of *A. lamprocarpa* and *A. wetarensis* were collected in this study because both species were distributed in NT and Wetar.

The geographic distribution of cpDNA haplotypes indicated that the NGI and QLD populations could be grouped into three regions, namely NGI, northern QLD, and southern QLD. There was a large difference in the cpDNA haplotypes among groups from different geographical areas; this was especially the case in the NGI and the QLD groups. This can be utilized to an improvement program as basic information for seed orchard establishment and provenance trial and so on. In general, the necessity of intensive seed collection from many populations should be emphasized; such collection should take place especially in southern QLD, where the highest genetic diversity is maintained.

V. CONCLUSION

Using Cold SSCP marker of 4 chloroplast DNA regions four cpDNA haplotypes were identified (A, B, C, and D). The haplotype distribution corresponded with the geographic distribution of the populations. Based on four cpDNA haplotypes, the eighteen populations were divided into three groups, namely, New Guinea Island, Northern Queensland and Southern Queensland.

ACKNOWLEDGEMENTS

The authors would like to thank JICA (Japan International Cooperation Agency) project for genetic materials (seeds) supporting. We would also like to thank Dr. Keiya Isoda for his technical assistance.

REFERENCES

- Balodis, V, and N. B. Clarkd. 1998. Tropical acacias - the new pulpwood. *Appita J.* 51, 179-181.
- Birky, C.W. 1988. Evolution and variation in plant chloroplast and mitochondrial genomes. In: Gottlieb, L. D. and Jain, S. K. (eds) *Plant Evolutionary Biology*, pp. 23-54. Chapman and Hall Ltd, London New York.
- Bodenes, C., F. Laigret, and A. Kremer. 1996. Inheritance and molecular variations of PCR-SSCP fragments in pedunculate oak (*Quercus robur L.*). *Theor. Appl. Genet.*, 93, 348-354.
- Bordáct, S., F. Popescu, D. Slade, U. M. Csaikl, I. Lesur, A. Borovics, P. Kézdy, A. O. König, D. Gömöry, S. Brewer, K. Burg and R. J. Petit. 2002. Chloroplast DNA variation of white oaks in northern Balkans and in the Carpathian Basin. *Forest Ecology and Management*, 156: 197-209.
- Brain, P. and B. R. Maslin. 1996. A serological investigation of the classification of *Acacia* subgenus Phyllodineae (Leguminosae:

- Mimosoideae). *Biochem. Sys. Eco.*, 24, 379-392.
- Brown, G. K., D. J. Murphy, J. T. Miller, and P. Y. Ladiges. 2008. *Acacia* s.s. and its Relationship Among Tropical Legumes, Tribe Ingeae (Leguminosae: Mimosoideae). *Systematic Botany* 33:739-751.
- Byrne, M., B. MacDonald, and D. Coates. 2002. Phylogeographical patterns in chloroplast DNA variation within the *Acacia acuminata* (Leguminosae: Mimosoideae) complex in Western Australia. *Journal of Evolutionary Biology* 15: 576-587.
- Byrne, M., G. Tischler, B. Macdonald, D. J. Coates, and J. McComb. 2001. Phylogenetic relationships between two rare acacias and their common, widespread relatives in south-western Australia. *Conservation Genetics* 2: 157-166.
- Clarke, H. D., S. R. Downie, and D. S. Seigler. 2000. Implications of chloroplast DNA restriction site variation for systematics of *Acacia* (Fabaceae: Mimosoideae). *Systematic Botany* 25: 618-632.
- Butcher, P., G. F. Moran, and H. D. Perkins. 1998. RFLP diversity in the nuclear genome of *Acacia mangium*. *Heredity*, 81, 205-213.
- Grace, M.B., G.S. Bambeck, G.S. Buzard, and B.D. Weintraub. 1995. Transverse temperature-gradient single-strand conformation polymorphism analysis for temperature optimization of 'Cold'-SSCP mutation detection. *Nucleic Acids. Res.* 23, 4224-4226.
- Guan L., Suharyanto, and S. Shiraishi. 2011. Isolation and characterization of tetranucleotide microsatellite loci in *Pinus massiniana* (Pinaceae). *Am. J. Bot.* 98: e216-e217.
- Hattori, K. 1997. Sequencing methods for PCR-amplified DNA. *Experimental Medicine* 15 (7), 711-717.
- Hayashi, K. 1991. PCR-SSCP: A simple and sensitive method for detection of mutation in genomic DNA. *PCR Methods Appl.*, 1, 34-38.
- Hong, Y-P., V. D. Hipkins, and S. H. Strauss. 1993. Chloroplast DNA diversity among trees, populations and species in the California closed-cone Pines (*Pinus radiata*, *Pinus muricata* and *Pinus attenuata*). *Genetics* 135: 1187-1196.
- Hongyo, T., G.S. Buzard, R.J. Calvert, and C. M. Weghorst. 1993. 'Cold SSCP' : a simple, rapid and non-radioactive method for optimized single-strand conformation polymorphism analysis. *Nucleic Acids Res.* 21: 3627-3642.
- Isoda, K., S. Shiraishi. 1999. Identification of chloroplast DNA haplotypes of *Abies firma* and *A. homolepis* using a polymerase chain reaction with species-specific primers.
- Judelson, H. S., L. J. Spielman, and R. C Shattock. 1995. Genetic-mapping and non-Mendelian of mating-type loci in the Oomycete, *Phytophthora infestans*. *Genetics* 141: 503-512.
- Khasa, P. D., W. M. Cheliak, and J. Bousquet. 1994. Genetic variation in 26 populations of *Racosperma auriculiforme* and *Racosperma mangium* using allozymes. *Can. J. For. Res.*, 24: 1123-1132.
- Levy, F., J. Antonovics, J. E. Boynton, and N. W. Gillham. 1996. A population genetic analysis of chloroplast DNA in *Phacelia*. *Heredity* 76: 143-155.
- Maeda, H., and S. Shiraishi. 1997. An identification of chloroplast DNA haplotypes of *Larix kaempferi* and *L. gmelinii* var. *japonica* using fluorescence-based PCR-SSCP analysis of *rbcl* gene. *J. For. Res.*, 2: 187-188.
- McDonald, M. W., and B. R. Maslin. 2000. Taxonomic Revision of the Salwoods: *Acacia aulacocarpa* Cunn. ex Benth. and its allies (Leguminosae: Mimosoideae: section Juliflorae). *Australian Systematic Botany* 13: 21-78.
- Moran, G. F., O. Muona, and J. C. Bell. 1988. *Acacia mangium*: A tropical forest tree of the coastal lowlands with low genetic diversity. *Evolution* 43: 231-235.
- Moran, G. F., O. Muona, and J. C. Bell. 1989. Breeding systems and genetic diversity in *Acacia auriculiformis* and *A. crassicarpa*. *Biotropica* 21: 250-256.

- Orita, M., H. Iwahana, H. Kanazawa, K. Hayashi, and T. Sekiya. 1989. Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphisms. *Proc. Natl. Acad. Sci. USA*, 86: 2766-2770.
- Pettigrew, C. J. and L. Watson. 1975. On the classification of Australian Acacias. *Aust. J. Bot.* 23: 833-847.
- Rehben, H., G. Kress, and T. Schmidt. 1997. Application of PCR-SSCP to species identification of fishery products. *J. Sci. Food Agric.* 74: 35-41.
- Spinardi, L., R. Mazars, and C. Theillet. 1991. Protocols for an improved detection of point mutations by SSCP. *Nucleic Acids Res.* 19: 4009.
- Taberlet, P., L. Gielly, G. Pautou, and J. Boubet. 1991. Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant Mol. Biol.* 17: 1105-1109.
- Thompson, L.A.J. 1994. *Acacia aulacocarpa*, *A. cincinnata*, *A. crassicarpa* and *A. wetarensis*: an annotated bibliography. Division of Forestry, Australian Tree Seed Centre, Canberra. 131pp.
- Watano, Y. M. Imazu and T. Shimizu. 1995. Chloroplast DNA typing by PCR-SSCP in the *Pinus pumila*-*P. parviflora* var. *pentaphylla* complex (Pinaceae). *J. Plant. Res.* 108: 493-499.
- Wickneswari, R. and M. Norwati. 1993. Genetic diversity of natural populations of *Acacia auriculiformis* in Australia and Papua New Guinea. *Aust. J. Bot.* 41: 65-77.
- Widyatmoko, AYPBC., A. Watanabe, and S. Shiraishi. 2010. Study on genetic variation and relationships among four *Acacia* species using RAPD and SSCP marker. *Journal of Forestry Research*, 7: 125-144.
- Widyatmoko, AYPBC. and S. Shiraishi. 2011. Sequence polymorphisms of four chloroplast genes in four *Acacia* species. *Journal of Forestry Research* 8: 79-89.
- Wolfe, K. H., W.-H. Li, and P. M. Sharp. 1987. Rates of nucleotide substitution vary greatly among plant mitochondrial, chloroplast, and nuclear DNAs. *Proc. Natl. Acad. Sci. USA* 84: 9054-9058.
- Zulfahmi, I. Z. Siregar, and U. J. Siregar. 2010. Chloroplast DNA variation of *Shorea acuminata* Dyer in Eastern Sumatra assessed by microsatellite markers. *Biodiversitas* 11 (3): 107-111.